

REFERENCES

1. Brandt CM, Sistrunk WW, Duffy MC *et al.* *Staphylococcus aureus* prosthetic joint infection treated with debridement and prosthesis retention. *Clin Infect Dis* 1997; **24**: 914–919.
2. Wilson MG, Kelley K, Thornhill TS. Infection as a complication of total knee-replacement arthroplasty. Risk factors and treatment in sixty-seven cases. *J Bone Joint Surg* 1990; **72**: 878–883.
3. Schoifet SD, Morrey BF. Treatment of infection after total knee arthroplasty by debridement with retention of the components. *J Bone Joint Surg Am* 1990; **72**: 1383–1390.
4. Drancourt M, Stein A, Argenson JN, Zannier A, Curvale G, Raoult D. Oral rifampin plus ofloxacin for treatment of *Staphylococcus*-infected orthopedic implants. *Antimicrob Agents Chemother* 1993; **37**: 1214–1218.
5. Javaloyas DM, Monreal PM. Oral antibiotic therapy in the adult bacterial osteomyelitis: results after two years of follow-up. *Med Clin (Barc)* 1999; **113**: 488–489.
6. Widmer AF, Gaechter A, Ochsner PE, Zimmerli W. Antimicrobial treatment of orthopedic implant-related infections with rifampin combinations. *Clin Infect Dis* 1992; **14**: 1251–1253.
7. Zimmerli W, Widmer AF, Blatter M, Frei R, Ochsner PE. Role of rifampin for treatment of orthopedic implant-related staphylococcal infections: a randomized controlled trial. Foreign-Body Infection (FBI) Study Group. *JAMA* 1998; **279**: 1537–1541.
8. Lewin CS, Amyes SG. The bactericidal activity of DR-3355, an optically active isomer of ofloxacin. *J Med Microbiol* 1989; **30**: 227–231.
9. Morris AB, Kanyok TP, Scott J *et al.* Rifamycins. In: Yu VL, Merigan TC, Barriere SL, eds, *Antimicrobial therapy and vaccines*. Philadelphia, PA: Lippencott, Williams & Wilkins, 1998; 901–962.
10. Stein A, Bataille JF, Drancourt M *et al.* Ambulatory treatment of multidrug-resistant *Staphylococcus*-infected orthopedic implants with high-dose oral co-trimoxazole (trimethoprim-sulfamethoxazole). *Antimicrob Agents Chemother* 1998; **42**: 3086–3091.
11. Bassetti M, Di Biagio A, Cenderello G *et al.* Linezolid treatment of prosthetic hip infections due to methicillin-resistant *Staphylococcus aureus* (MRSA). *J Infect* 2001; **43**: 148–149.
12. Rayner CR, Baddour LM, Birmingham MC, Norden C, Meagher AK, Schentag JJ. Linezolid in the treatment of osteomyelitis: results of compassionate use experience. *Infection* 2004; **32**: 8–14.
13. Bassetti M, Vitale F, Melica G *et al.* Linezolid in the treatment of Gram-positive prosthetic joint infections. *J Antimicrob Chemother* 2005; **55**: 387–390.
14. Deirmengian C, Greenbaum J, Lotke PA, Booth RE, Lonner JH. Limited success with open debridement and retention of components in the treatment of acute *Staphylococcus aureus* infections after total knee arthroplasty. *J Arthroplasty* 2003; **18** (suppl 1): 22–26.

RESEARCH NOTE

Clinical significance of isolated *Staphylococcus aureus* central venous catheter tip cultures

J. J. Ruhe and A. Menon

Division of Infectious Diseases, Department of Medicine, University of Arkansas for Medical Sciences and the Central Arkansas Veterans Healthcare System, Little Rock, AR, USA

ABSTRACT

This retrospective cohort study examined the clinical significance of isolated *Staphylococcus aureus* central venous catheter (CVC) tip cultures (i.e., positive tip cultures without concomitant positive blood cultures). Subsequent *S. aureus* bacteraemia was found in nine (12%) of 77 patients at a median time of 4 days after CVC removal. A high co-morbidity score and no effective antibiotic treatment within 48 h of CVC removal were independent risk-factors for septic complications following multivariate analysis. A matched case-control study that compared the above cohort with patients with CVC tip cultures negative for *S. aureus* supported the significance of these findings.

Keywords Bacteraemia, central venous catheter, risk-factors, significance, *Staphylococcus aureus*, tip cultures

Original Submission: 28 December 2005; **Revised Submission:** 13 February 2006; **Accepted:** 1 March 2006

Clin Microbiol Infect 2006; **12**: 933–936
10.1111/j.1469-0691.2006.01491.x

Few data exist concerning the clinical significance of central venous catheter (CVC) tip cultures that are positive for *Staphylococcus aureus* in patients who have no blood cultures collected around the time of CVC removal, or whose

Corresponding author and reprint requests: J. J. Ruhe, Division of Infectious Diseases, University of Arkansas for Medical Sciences, 4301 West Markham Street, Slot #639, Little Rock, AR 72205, USA
E-mail: ruhejorgj@uams.edu

concomitant blood cultures do not grow *S. aureus* (i.e., isolated CVC tip cultures) [1]. The present report describes a retrospective cohort study of all adult patients with isolated *S. aureus* CVC tip cultures at two tertiary care medical centres between January 1999 and December 2004.

Cases were defined as patients who: (i) had a distal CVC tip culture that grew ≥ 15 *S. aureus* colonies by the roll-plate technique, and (ii) either did not have blood cultures taken, or had negative blood cultures taken between 2 days before and 1 day after CVC removal (concomitant blood cultures) [2]. A subsequent septic complication was defined as isolation of *S. aureus* with an antibiotic susceptibility pattern identical to the CVC tip isolate from a normally sterile site between 3 days and 12 weeks after CVC removal. Additionally, a matched, nested, 1:1 case-control study was performed to compare the *S. aureus* complication rate in the cases with that in a similar patient cohort with CVC tip cultures negative for *S. aureus*. Control patients were selected randomly and matched by age, Charlson index, and stay in a surgical or medical intensive care unit. The Charlson index reflects the sum of a patient's weighted co-morbidities and predicts mortality [3–5]. Therapy was considered appropriate if treatment with at least one antibiotic to which susceptibility was documented was started within 24 h before and 48 h after CVC removal (adapted from [6]). *S. aureus* processing was performed according to CLSI guidelines [7,8]. The study was approved by the institutional review boards of the hospitals concerned.

Septic complication was the outcome of interest. Bivariate analyses were conducted by Pearson's chi-square test, Fisher's exact test or the Mann–Whitney *U*-test, as appropriate, to identify factors associated with the outcome ($p < 0.05$). Variables with $p < 0.2$ were entered into a backward stepwise non-conditional logistic regression analysis to identify independent risk-factors. In the case-control study, McNemar's test and the Wilcoxon signed-rank test were used to compare categorical and continuous variables, respectively; the log-rank test was used to compare the rates of septic complications.

Of 101 isolated *S. aureus* CVC tip cultures, 24 were excluded for the following reasons: follow-up of ≤ 12 weeks or incomplete medical records ($n = 11$); invasive *S. aureus* infection in ≤ 12 weeks

($n = 3$); death within 48 h of CVC removal ($n = 3$); CVC tip with multiple organisms ($n = 3$); concurrent non-bacteraemic *S. aureus* infection ($n = 2$); and *S. aureus* bacteraemia 24–48 h after CVC removal ($n = 2$). The final cohort comprised 77 consecutive patients with a mean age of 59 years; nine (12%) of 77 patients developed a subsequent *S. aureus*-related complication, all of which were bacteraemia. One patient died from *S. aureus*-related septic shock.

Table 1 shows the results of the bivariate analyses. A high Charlson index, chronic obstructive pulmonary disease (COPD), and no effective antibiotic therapy received within 48 h of CVC removal, were all factors associated with septic complications. Only two (3%) of 59 patients who had received appropriate antibiotics within 48 h of CVC removal developed subsequent bacteraemia, compared with seven (39%) of 18 patients who had not received appropriate antibiotics. The median duration of antibiotic treatment in patients with no complication was 12 days.

Table 1. Risk-factors for *Staphylococcus aureus*-related septic complications among 77 patients with isolated *S. aureus* central venous catheter tip cultures

Characteristic	Cases with complications ^a (<i>n</i> = 9)	Cases without complications ^a (<i>n</i> = 68)	OR (95% CI)	<i>p</i>
Mean age \pm SD (years)	55 \pm 24	59 \pm 17		> 0.2
Male gender	7 (78)	44 (65)	1.91 (0.37–9.92)	> 0.2
Stay in ICU	3 (33)	26 (38)	0.81 (0.19–3.51)	> 0.2
Co-morbid conditions				
Charlson score ≥ 5	5 (56)	10 (15)	7.25 (1.66–31.71)	0.01
Diabetes mellitus	5 (56)	17 (25)	3.75 (0.90–15.59)	0.11
COPD	5 (56)	10 (15)	7.25 (1.66–31.71)	0.01
Haemodialysis dependency	0 (0)	3 (4)	–	> 0.2
Current malignancy	3 (33)	20 (29)	1.20 (0.27–5.28)	> 0.2
Intravascular device ^b	2 (22)	12 (18)	1.33 (0.25–7.23)	> 0.2
Orthopaedic device	0 (0)	10 (15)	–	> 0.2
Previous surgery ^c	4 (44)	34 (50)	0.80 (0.20–3.24)	> 0.2
CVC exit site infection	3 (33)	28 (41)	0.71 (0.17–3.10)	> 0.2
SIRS ^d	4 (44)	39 (57)	0.59 (0.15–2.41)	> 0.2
Non-tunnelled CVC ^e	7 (78)	54 (79)	0.91 (0.17–4.86)	> 0.2
Concomitant blood cultures performed	4 (44)	40 (59)	0.56 (0.14–2.27)	> 0.2
MRSA	6 (67)	41 (60)	1.32 (0.30–5.72)	> 0.2
No effective antibiotics by day 2	7 (78)	11 (16)	18.14 (3.32–99.16)	< 0.001
28-day mortality rate	1 (11)	6 (9)	1.29 (0.14–12.15)	> 0.2

ICU, intensive care unit; COPD, chronic obstructive pulmonary disease; CVC, central venous catheter; SIRS, systemic inflammatory response syndrome; MRSA, methicillin-resistant *Staphylococcus aureus*.

^aTotal number (percentage) or mean \pm SD.

^bInferior vena cava filter ($n = 5$), pacemaker or implantable cardiac defibrillator ($n = 3$), synthetic intravascular graft ($n = 2$), prosthetic heart valve ($n = 2$), coronary artery stent ($n = 1$), transjugular intrahepatic portosystemic shunt ($n = 1$).

^cWithin the previous 6 months.

^dSee [17].

^eOther: tunnelled CVCs ($n = 7$), totally implantable devices ($n = 7$), and peripherally inserted central catheters ($n = 2$).

Table 2. Comparison of patient characteristics in the matched nested case-control study

Characteristic	Cases (%) (n = 77)	Controls (%) (n = 77)	P
Mean age \pm SD (years) ^a	59 \pm 18	59 \pm 15	> 0.2
Male gender	51 (66)	47 (61)	> 0.2
Stay in ICU ^a	29 (38)	29 (38)	> 0.2
Co-morbid conditions			
Median Charlson score (range) ^a	2 (0–11)	2 (0–10)	> 0.2
Diabetes mellitus	22 (29)	16 (21)	> 0.2
COPD	15 (20)	19 (25)	> 0.2
Chronic liver disease	8 (10)	8 (10)	> 0.2
Haemodialysis dependence	3 (4)	2 (3)	> 0.2
Current malignancy	23 (30)	29 (38)	> 0.2
Intravascular device ^b	14 (18)	12 (16)	> 0.2
Orthopaedic device	10 (13)	3 (4)	0.15
CVC exit site infection	31 (40)	17 (22)	0.06
SIRS	43 (56)	40 (52)	> 0.2
Non-tunnelled CVC	61 (79)	63 (82)	> 0.2
Days of hospitalisation after CVC tip culture obtained (range)	9 (0–82)	14 (0–122)	> 0.2
Complicating <i>Staphylococcus aureus</i> bacteraemia ^c	9 (12)	2 (2)	0.03
28-day mortality rate	7 (9)	11 (14)	> 0.2

ICU, intensive care unit; COPD, chronic obstructive pulmonary disease; CVC, central venous catheter; SIRS, systemic inflammatory response syndrome.

^aThese variables were used in the matching process.

^bInferior vena cava filter (*n* = 8), pacemaker or implantable cardiac defibrillator (*n* = 7), synthetic intravascular graft (*n* = 5), prosthetic heart valve (*n* = 3), intravascular stent (*n* = 2), transjugular intrahepatic portosystemic shunt (*n* = 1).

^c≤ 12 weeks.

Logistic regression analysis revealed two independent risk-factors for septic complications, namely a Charlson index ≥ 5 (adjusted OR 9.07; 95% CI 1.42–57.92; *p* 0.02) and no effective antibiotic therapy received within 48 h of CVC removal (adjusted OR 21.23; 95% CI 3.20–140.77; *p* 0.002).

Table 2 shows the results of the case-control study. Only two (2%) control patients, compared with nine (12%) case patients, developed septic complications (*p* 0.03); these occurred at a median of 62 and 4 days, respectively, after CVC removal (*p* 0.03).

S. aureus catheter tip cultures are accompanied frequently by positive blood cultures [9,10], and findings at the institutions in the present study, in which 267 (73%) of 368 *S. aureus* CVC tip cultures were associated with concurrent bacteraemia, agree with this observation. However, the clinical significance of isolated *S. aureus* CVC tip cultures has not been defined [1]. To our knowledge, the present study is the first to investigate specifically the outcome of patients with this clinical entity. Nine (12%) patients with isolated *S. aureus* CVC tip cultures were identified who developed bacteraemia between 3 and 33 days after catheter removal. A high Charlson index and a failure to

receive effective antibiotics within 48 h of CVC removal were the only independent risk-factors for subsequent septic complications. The latter also remained the only independent predictor in an analysis of a subgroup of 44 patients who had negative concomitant blood cultures (data not shown). This finding, together with the results of the matched case-control study, supports the conclusion that the subsequent bacteraemic episodes signify a genuine complication, rather than merely non-detection of *S. aureus* bacteraemia at the time of CVC removal or the existence of an independent healthcare-related complication.

COPD has not been associated with *S. aureus* bacteraemia in previous reports; its significance in bivariate analysis may be an artefactual finding caused by multiple testing. The majority (57%) of non-tunnelled CVCs was inserted into the subclavian vein. Catheter location and rate of septic complications were not related (*p* > 0.2). The pathogenic mechanisms responsible for subsequent *S. aureus* septic complications are unknown. The potential role of small, clinically non-evident, intravascular thrombi, or low-grade bacteraemia caused by mechanical manipulation at the time of CVC removal, remain to be determined [11].

There were limitations to the study. Blood and CVC tip isolates were not available for molecular typing [12,13], so their relatedness could not be confirmed. Only 21% of the patients underwent echocardiography; therefore, endocarditis may have been under-diagnosed [14,15]. The study was also limited by its retrospective design; however, strict inclusion criteria were used to minimise sampling, misclassification and surveillance bias [16].

In summary, isolated *S. aureus* CVC tip cultures were associated with a significant risk of septic complications. If validated, the data would indicate that patients with this clinical entity should receive a course of anti-staphylococcal antibiotics, the optimal duration of which remains to be determined.

ACKNOWLEDGEMENTS

This study was presented, in part, at the 42nd Annual Meeting of the Infectious Diseases Society of America (Boston, 2004). The authors would like to thank L. Myers for statistical advice, and G. Schutze for his review of the manuscript.

REFERENCES

1. Mermel LA, Farr BM, Sherertz RJ, Raad II, O'Grady N, Harris JS. Guidelines for the management of intravascular catheter-related infections. *Clin Infect Dis* 2001; **32**: 1249–1272.
2. Maki DG, Weise CE, Sarafin HW. A semiquantitative culture method for identifying intravenous catheter-related infection. *N Engl J Med* 1977; **296**: 1305–1309.
3. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987; **40**: 373–383.
4. Lesens O, Methlin C, Hansmann Y *et al.* Role of comorbidity in mortality related to *Staphylococcus aureus* bacteremia: a prospective study using the Charlson weighted index of comorbidity. *Infect Control Hosp Epidemiol* 2003; **24**: 890–896.
5. Beddhu S, Bruns FJ, Saul M *et al.* A simple comorbidity scale predicts clinical outcomes and costs in dialysis patients. *Am J Med* 2000; **108**: 609–613.
6. Fowler VG, Olsen MK, Corey R *et al.* Clinical identifiers of complicated *Staphylococcus aureus* bacteremia. *Arch Int Med* 2003; **163**: 2066–2072.
7. NCCLS. *Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically*, 6th edn. Approved standard M7-A6. Wayne, PA: NCCLS, 2003.
8. NCCLS. *Performance standards for antimicrobial susceptibility testing*. 13th informational supplement M100–S13. Wayne, PA: NCCLS, 2003.
9. Cercenado E, Ena J, Rodriguez-Creixems M, Romero I, Bouza E. A conservative procedure for the diagnosis of catheter-related infections. *Arch Intern Med* 1990; **150**: 1417–1420.
10. Peacock SJ, Eddleston M, Emptage A, King A, Crook DW. Positive intravenous line tip cultures as predictors of bacteraemia. *J Hosp Infect* 1998; **40**: 35–38.
11. Raad I, Narro J, Khan A, Tarrand J, Vartivarian S, Bodey GP. Serious complications of vascular catheter-related *Staphylococcus aureus* bacteremia in cancer patients. *Eur J Clin Microbiol Infect Dis* 1992; **11**: 675–682.
12. Tenover FC, Arbeit RD, Goering RV *et al.* 1995. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol* 1995; **33**: 2233–2239.
13. Khatib R, Sharma M, Naqvi SA *et al.* Molecular analysis of *Staphylococcus aureus* blood isolates shows lack of polyclonal bacteremia. *J Clin Microbiol* 2003; **41**: 1717–1719.
14. Fätkenheuer G, Cornely O, Seifert H. Clinical management of catheter-related infections. *Clin Microbiol Infect* 2002; **8**: 545–550.
15. Fowler VG, Li J, Corey GR *et al.* Role of echocardiography in evaluation of patients with *Staphylococcus aureus* bacteremia: experience in 103 patients. *J Am Coll Cardiol* 1997; **30**: 1072–1078.
16. Jernigan JA, Farr BM. Short-course therapy of catheter-related *Staphylococcus aureus* bacteremia: a meta-analysis. *Ann Intern Med* 1993; **119**: 304–311.
17. American College of Chest Physicians/Society of Critical Care Medicine Consensus Conference Committee. Definition for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Crit Care Med* 1992; **20**: 864–874.

RESEARCH NOTE

Evaluation of differential gene expression in susceptible and resistant clinical isolates of *Klebsiella pneumoniae* by DNA microarray analysis

A. Doménech-Sánchez¹, V. Javier Benedí¹,
L. Martínez-Martínez² and S. Albertí¹

¹Departamento de Biología and Institut Universitari d'Investigació en Ciències de la Salut (IUNICS), Universidad de las Islas Baleares, Palma de Mallorca and ²Servicio de Microbiología, Hospital Universitario Marqués de Valdecilla, Santander, Spain

ABSTRACT

DNA microarray technology was used to evaluate differential gene expression in a susceptible *Klebsiella pneumoniae* isolate and a resistant clinical derivative. Nineteen genes were up-regulated in the resistant isolate when compared with the susceptible isolate. An ABC transporter-related gene, *ycjV*, was strongly over-expressed, suggesting the existence of a novel active efflux mechanism. Approximately half of the up-regulated genes coded for ribosomal proteins, or proteins involved in tRNA metabolism. Among 33 down-regulated genes, almost one-third were related to nitrogen metabolism. A possible role of fitness in the development of antimicrobial resistance is suggested.

Keywords Antimicrobial resistance, efflux pumps, gene regulation, *Klebsiella pneumoniae*, microarray, transcriptional analysis

Original Submission: 21 June 2005; **Revised Submission:** 25 January 2006; **Accepted:** 6 February 2006

Clin Microbiol Infect 2006; **12**: 936–940
10.1111/j.1469-0691.2006.01470.x

Corresponding author and reprint requests: S. Albertí, IUNICS, Edificio Guillen Colom, CAMPUS-UIB, Crte. Valldemorsa, km 7.5 Palma de Mallorca 07122, Spain
E-mail: sabastian.alberti@uib.es